

# The largest Hosoya index of $(n, n + 1)$ -graphs<sup>☆</sup>

Hanyuan Deng

College of Mathematics and Computer Science, Hunan Normal University, Changsha, Hunan 410081, PR China

## ARTICLE INFO

### Article history:

Received 18 July 2007

Received in revised form 29 April 2008

Accepted 27 May 2008

### Keywords:

$(n, n + 1)$ -graph

Hosoya index

Fibonacci number

Matching

Transformation

## ABSTRACT

The Hosoya index of a graph is defined as the total number of its matchings. In this paper, we obtain that the largest Hosoya index of  $(n, n + 1)$ -graphs is  $f(n + 1) + f(n - 1) + 2f(n - 3)$ , where  $f(n)$  is the  $n$ th Fibonacci number, and we characterize the extremal graphs.

© 2008 Elsevier Ltd. All rights reserved.

## 1. Introduction

The Hosoya index or Z-index  $z(G)$  of a graph  $G$  is the total number of its matchings plus one; the one corresponds to a matching in a set with zero edges. The Hosoya index is a prominent example of topological indices which are of interest in combinatorial chemistry. It is defined as the total number of matchings (independent edge subsets) of a graph. The Z-index was introduced by Hosoya [1,2] in 1971, and it turned out to be applicable to several questions of molecular chemistry. For example, the connections with physico-chemical properties such as boiling point, entropy or heat of vaporization are well studied.

Several papers deal with the characterization of the extremal graphs with respect to the Hosoya index in several given graph classes. Usually, trees, unicyclic graphs and certain structures involving pentagonal and hexagonal cycles are of major interest [3–12].

In this paper, we deal with the characterization of the extremal graphs with the largest Hosoya index among all  $(n, n + 1)$ -graphs, i.e., the simple connected graphs with  $n$  vertices and  $n + 1$  edges.

Let  $G = (V, E)$  be a simple connected graph with the vertex set  $V$  and the edge set  $E$ . For any  $v \in V$ ,  $N_G(v)$  denotes the neighbors of  $v$ , and  $d_G(v) = |N_G(v)|$  is the degree of  $v$ . A leaf is a vertex of degree one.

If  $E' \subseteq E(G)$  and  $W \subseteq V(G)$ , then  $G - E'$  and  $G - W$  denote the subgraphs of  $G$  obtained by deleting the edges of  $E'$  and the vertices of  $W$ , respectively. If a graph  $G$  has components  $G_1, G_2, \dots, G_t$ , then  $G$  is denoted by  $\bigcup_{i=1}^t G_i$ . We denote by  $P_n$  the path on  $n$  vertices and by  $C_n$  the cycle on  $n$  vertices.

The following basic results will be used.

(i) If  $v$  is a vertex of  $G$ , then

$$z(G) = z(G - \{v\}) + \sum_{x \in N_G(v)} z(G - \{v, x\}). \quad (1)$$

<sup>☆</sup> Project 10471037 supported by the National Natural Science Foundation of China.

E-mail address: [hydeng@hunnu.edu.cn](mailto:hydeng@hunnu.edu.cn).




Fig. 1. The subgraphs induced by the cycles of an  $(n, n + 1)$ -graph.




Fig. 2. Transformation 1.




Fig. 3. The graphs in Remark 1.

(ii) If  $uv$  is an edge of  $G$ , then

$$z(G) = z(G - uv) + z(G - \{u, v\}). \quad (2)$$

(iii) If  $G$  is a graph with components  $G_1, G_2, \dots, G_k$ , then

$$z(G) = \prod_{i=1}^k z(G_i).$$

(vi)  $z(C_n) = f(n - 1) + f(n + 1)$  for  $n \geq 3$ .

(v)  $z(P_0) = 1, z(P_1) = 1$  and  $z(P_n) = f(n + 1)$  for  $n \geq 2$

where  $f(0) = 0, f(1) = 1$  and  $f(n) = f(n - 1) + f(n - 2)$  for  $n \geq 2$  denotes the sequence of Fibonacci numbers.

The following property of the Fibonacci numbers is very well known and can be seen immediately by plugging into Binet's formula.

**Lemma 1.1.**  $f(n) = f(k)f(n - k + 1) + f(k - 1)f(n - k)$  for  $1 \leq k \leq n$ .

Let  $\mathcal{G}(n, n + 1)$  be the set of simple connected graphs with  $n$  vertices and  $n + 1$  edges. For any graph  $G \in \mathcal{G}(n, n + 1)$ , there are two cycles  $C_p$  and  $C_q$  in  $G$ . We divide all the  $(n, n + 1)$ -graphs with two cycles of lengths  $p$  and  $q$  into three classes.

(1)  $\mathcal{A}(p, q)$  is the set of  $G \in \mathcal{G}(n, n + 1)$  in which the cycles  $C_p$  and  $C_q$  have only one common vertex;

(2)  $\mathcal{B}(p, q)$  is the set of  $G \in \mathcal{G}(n, n + 1)$  in which the cycles  $C_p$  and  $C_q$  have no common vertex;

(3)  $\mathcal{C}(p, q, r)$  is the set of  $G \in \mathcal{G}(n, n + 1)$  in which the cycles  $C_p$  and  $C_q$  have a common path of length  $r + 1, r \geq 0$ .

Note that the subgraph induced by the cycles of  $G \in \mathcal{A}(p, q)$  (or  $\mathcal{B}(p, q)$ ,  $\mathcal{C}(p, q, r)$ ) is shown in Fig. 1(a) (or (b),(c)) and  $\mathcal{C}(p, q, r) = \mathcal{C}(p, p + q - 2r + 2, p - r - 2) = \mathcal{C}(p + q - 2r - 2, q, q - r - 2)$ .

## 2. Transformations increasing the Hosoya index

For convenience, we introduce in this section three transformations, which increase the Hosoya index.

**Transformation 1.** Let  $G \neq P_1$  be a connected graph and choose  $u \in V(G)$ .  $G_1$  denotes the graph that results from identifying  $u$  with the vertex  $v_k$  of a simple path  $v_1 v_2 \dots v_n, 1 < k < n$ ;  $G_2$  is obtained from  $G_1$  by deleting  $v_{k-1} v_k$  and adding  $v_1 v_n$  (see Fig. 2).

**Lemma 2.1** ([9]). Let  $G_1$  and  $G_2$  be the graphs in Fig. 2. Then  $z(G_1) < z(G_2)$ .

**Remark 1.** Repeating Transformation 1, any tree  $T$  of size  $t$  attached to a graph  $G$  can be changed into a path with  $t$  edges as shown in Fig. 3. The Hosoya index increases by this transformation.

**Transformation 2.** Let  $P = uu_1 u_2 \dots u_t v$  be a path in  $G$  and  $G \neq P$ , where the degrees of  $u_1, \dots, u_t$  in  $G$  are 2 (such a path  $P$  is called an internal path of  $G$ ).  $G_1$  denotes the graph that results from identifying  $u$  with the vertex  $v_k$  of a simple path  $v_1 v_2 \dots v_k$  and identifying  $v$  with the vertex  $v_{k+1}$  of another simple path  $v_{k+1} v_{k+2} \dots v_n, 1 < k < n - 1$ ;  $G_2$  is obtained from  $G_1$  by deleting  $v_{k-1} v_k$  and adding  $v_1 v_n$ ,  $G_3$  is obtained from  $G_1$  by deleting  $v_{k+1} v_{k+2}$  and adding  $v_1 v_n$  (see Fig. 4).




Fig. 4. Transformation 2.

**Lemma 2.2.** Let  $G_1$ ,  $G_2$  and  $G_3$  be the graphs in Fig. 4. Then  $z(G_1) < z(G_2)$  or  $z(G_1) < z(G_3)$ .

**Proof.** Let  $X = N_G(u) - \{u_1\}$  and  $Y = N_G(v) - \{u_t\}$ .

$$A = z(G - \{u, u_1, \dots, u_t, v\})$$

$$B = \sum_{y \in Y} z(G - \{u, u_1, \dots, u_t, v, y\})$$

$$C = \sum_{x \in X} z(G - \{u, u_1, \dots, u_t, v, x\})$$

$$D = \sum_{x \in X} \sum_{y \in Y} z(G - \{u, u_1, \dots, u_t, v, x, y\}).$$

By Eq. (1),

$$\begin{aligned} z(G_1) &= z(G_1 - \{u\}) + \sum_{x \in X} z(G_1 - \{u, x\}) + z(G_1 - \{u, u_1\}) + z(G_1 - \{u, v_{k-1}\}) \\ &= z(G_1 - \{u, v\}) + \sum_{y \in Y} z(G_1 - \{u, v, y\}) + z(G_1 - \{u, v, u_t\}) + z(G_1 - \{u, v, v_{k+2}\}) \\ &\quad + \sum_{x \in X} z(G_1 - \{u, x, v\}) + \sum_{x \in X} \sum_{y \in Y} z(G_1 - \{u, x, v, y\}) + \sum_{x \in X} z(G_1 - \{u, x, v, u_t\}) \\ &\quad + \sum_{x \in X} z(G_1 - \{u, x, v, v_{k+2}\}) + z(G_1 - \{u, u_1, v\}) + \sum_{y \in Y} z(G_1 - \{u, u_1, v, y\}) + z(G_1 - \{u, u_1, v, u_t\}) \\ &\quad + z(G_1 - \{u, u_1, v, v_{k+2}\}) + z(G_1 - \{u, v_{k-1}, v\}) + \sum_{y \in Y} z(G_1 - \{u, v_{k-1}, v, y\}) \\ &\quad + z(G_1 - \{u, v_{k-1}, v, u_t\}) + z(G_1 - \{u, v_{k-1}, v, v_{k+2}\}) \\ &= Af(k)f(n-k)f(t+1) + Bf(k)f(n-k)f(t+1) + Af(k)f(n-k)f(t) \\ &\quad + Af(k)f(n-k-1)f(t+1) + Cf(k)f(n-k)f(t+1) + Df(k)f(n-k)f(t+1) \\ &\quad + Cf(k)f(n-k)f(t) + Cf(k)f(n-k-1)f(t+1) + Af(k)f(n-k)f(t) \\ &\quad + Bf(k)f(n-k)f(t) + Af(k)f(n-k)f(t-1) + Af(k)f(n-k-1)f(t) \\ &\quad + Af(k-1)f(n-k)f(t+1) + Bf(k-1)f(n-k)f(t+1) \\ &\quad + Af(k-1)f(n-k)f(t) + Af(k-1)f(n-k-1)f(t+1) \\ &= A[f(k)f(n-k)f(t+1) + f(k)f(n-k)f(t) + f(k)f(n-k-1)f(t+1) \\ &\quad + f(k)f(n-k)f(t) + f(k)f(n-k)f(t-1) + f(k)f(n-k-1)f(t) \\ &\quad + f(k-1)f(n-k)f(t+1) + f(k-1)f(n-k)f(t) + f(k-1)f(n-k-1)f(t+1)] \\ &\quad + B[f(k)f(n-k)f(t+1) + f(k)f(n-k)f(t) + f(k-1)f(n-k)f(t+1)] \\ &\quad + C[f(k)f(n-k)f(t+1) + f(k)f(n-k)f(t) + f(k)f(n-k-1)f(t+1)] \\ &\quad + Df(k)f(n-k)f(t+1) \\ &= A[f(k+1)f(n-k+1)f(t+1) + f(k+1)f(n-k)f(t) + f(k)f(n-k+1)f(t) \\ &\quad + f(k)f(n-k)f(t-1)] + B[f(k+1)f(n-k)f(t+1) + f(k)f(n-k)f(t)] \\ &\quad + C[f(k)f(n-k+1)f(t+1) + f(k)f(n-k)f(t)] \\ &\quad + Df(k)f(n-k)f(t+1). \end{aligned}$$




Fig. 5. The graphs  $H_i(p, q)$ ,  $i = 1, 2, 3$ .

Similarly, we have

$$z(G_2) = A[f(n)f(t+1) + f(n+1)f(t) + f(n-1)f(t-1)] + B[f(n-1)f(t+1) + f(n-1)f(t)] \\ + C[f(n)f(t+1) + f(n-1)f(t)] + Df(n-1)f(t+1);$$

$$z(G_3) = A[f(n)f(t+1) + f(n+1)f(t) + f(n-1)f(t-1)] + B[f(n)f(t+1) + f(n-1)f(t)] \\ + C[f(n-1)f(t+1) + f(n-1)f(t)] + Df(n-1)f(t+1).$$

If  $B \leq C$ , then by Lemma 1.1

$$\begin{aligned} \Delta_1 &= z(G_2) - z(G_1) \\ &= A[(f(n) - f(k+1)f(n-k+1))f(t+1) + (f(n+1) - f(k+1)f(n-k) \\ &\quad - f(k)f(n-k+1))f(t) + (f(n-1) - f(k)f(n-k))f(t-1)] \\ &\quad + B[(f(n-1) - f(k+1)f(n-k))f(t+1) + (f(n-1) - f(k)f(n-k))f(t)] \\ &\quad + C[(f(n) - f(k)f(n-k+1))f(t+1) + (f(n-1) - f(k)f(n-k))f(t)] \\ &\quad + D[f(n-1) - f(k)f(n-k)]f(t+1) \\ &= A[-f(k-1)f(n-k-1)f(t+1) + f(k-1)f(n-k-1)f(t) + f(k-1)f(n-k-1)f(t-1)] \\ &\quad + B[-f(k-1)f(n-k-2)f(t+1) + f(k-1)f(n-k-1)f(t)] \\ &\quad + C[f(k-1)f(n-k)f(t+1) + f(k-1)f(n-k-1)f(t)] + Df(k-1)f(n-k-1)f(t+1) \\ &= B[-f(k-1)f(n-k-2)f(t+1) + f(k-1)f(n-k-1)f(t)] \\ &\quad + C[f(k-1)f(n-k)f(t+1) + f(k-1)f(n-k-1)f(t)] + Df(k-1)f(n-k-1)f(t+1) > 0. \end{aligned}$$

If  $B > C$ , then by Lemma 1.1

$$\begin{aligned} \Delta_2 &= z(G_3) - z(G_1) \\ &= A[(f(n) - f(k+1)f(n-k+1))f(t+1) + (f(n+1) - f(k+1)f(n-k) \\ &\quad - f(k)f(n-k+1))f(t) + (f(n-1) - f(k)f(n-k))f(t-1)] \\ &\quad + B[(f(n) - f(k+1)f(n-k))f(t+1) + (f(n-1) - f(k)f(n-k))f(t)] \\ &\quad + C[(f(n-1) - f(k)f(n-k+1))f(t+1) + (f(n-1) - f(k)f(n-k))f(t)] \\ &\quad + D[f(n-1) - f(k)f(n-k)]f(t+1) \\ &= B[f(k)f(n-k-1)f(t+1) + f(k-1)f(n-k-1)f(t)] \\ &\quad + C[-f(k-2)f(n-k-1)f(t+1) + f(k-1)f(n-k-1)f(t)] + Df(k-1)f(n-k-1)f(t+1) > 0. \end{aligned}$$

The proof is completed.  $\square$

**Remark 2.** If  $G$  is the  $(n, n+1)$ -graph with trees  $T_1, \dots, T_k$  of sizes  $t_1, \dots, t_k$  attached to one of the graphs  $H_i(p, q)$  ( $i = 1, 2, 3$ ) in Fig. 5, then these trees are changed into paths of sizes  $t_1, \dots, t_k$  after repeating Transformation 1. Let us continue repeating Transformation 2, these paths will be combined into one path. So,  $G$  can be changed into an  $(n, n+1)$ -graph with at least one path attached to one of the graphs  $H_i(p, q)$ , and the Hosoya index increases by these transformations.

The proof of the following lemma is obvious.

**Lemma 2.3.** (i) Let  $P = v_1 v_2 \dots v_n$  be a path of length  $n-1$ ,  $n > 1$ .  $M_1(P)$  is the set of matchings of  $P$  containing  $v_1 v_2$  and  $M_2(P)$  is the set of matchings of  $P$  containing  $v_n v_{n-1}$ . Define  $\xi_P(v_i v_{i+1}) = v_{n-i+1} v_{n-i}$  for  $i = 1, 2, \dots, n-1$ . Then  $\xi_P$  is a bijective mapping between  $M_2(P)$  and  $M_1(P)$ ;

(ii) Let  $C = v_1 v_2 \dots v_n v_{n+1}$  be a cycle of length  $n$ ,  $v_{n+1} = v_1$ .  $M_1(C)$  is the set of matchings of  $C$  containing  $v_1 v_2$  and  $M_2(C)$  is the set of matchings of  $C$  containing  $v_1 v_n$ .  $\xi_C(v_i v_{i+1}) = v_{n-i+2} v_{n-i+1}$  for  $i = 1, 2, \dots, n$ . Then  $\xi_C$  is a bijective mapping between  $M_2(C)$  and  $M_1(C)$ .

**Transformation 3.** Let  $P = u_0 u_1 u_2 \dots u_{t+1}$  be a path or a cycle (if  $u_0 = u_{t+1}$ ) in  $G$ , where the degrees of  $u_1, \dots, u_t$  in  $G$  are 2,  $t \geq 1$ .  $G_1$  denotes the graph that results from identifying  $u_r$  ( $0 \leq r \leq t$ ) with the vertex  $v_k$  of a simple path  $v_1 v_2 \dots v_k$ ;  $G_2$  is obtained from  $G_1$  by deleting  $u_r u_{r+1}$  and adding  $u_{r+1} v_1$  (see Fig. 6).




Fig. 6. Transformation 3.

**Lemma 2.4.** Let  $G_1$  and  $G_2$  be the graphs in Fig. 6. Then  $z(G_1) < z(G_2)$ .

**Proof.** Let  $M(G_1)$  and  $M(G_2)$  be the sets of matchings of  $G_1$  and  $G_2$ , respectively. We construct a mapping  $\eta : M(G_1) \rightarrow M(G_2)$ ,  $\forall B \in M(G_1)$

$$\eta(B) = \begin{cases} (B - \{u_r u_{r+1}\} - B \cap P^*) \cup \{u_{r+1} v_1\} \cup \xi_{P^*}(B \cap P^*), & \text{if } u_r u_{r+1} \in B \text{ and } v_1 v_2 \in B; \\ (B - \{u_r u_{r+1}\}) \cup \{u_{r+1} v_1\}, & \text{if } u_r u_{r+1} \in B \text{ and } v_1 v_2 \notin B; \\ B, & \text{otherwise} \end{cases}$$

where  $P^* = u_r v_{k-1} \cdots v_2 v_1$  and  $\xi_{P^*}$  is defined in Lemma 2.3.

Note that  $u_r v_{k-1} \notin B$  if  $u_r u_{r+1} \in B$ . The mapping  $\eta$  is injective. However, there is no  $B \in M(G_1)$  with  $\eta(B) = \{u_r u_{r-1}, u_{r+1} v_1\}$ . So,  $z(G_1) < z(G_2)$ .

The proof is completed.  $\square$

**Remark 3.** Repeating Transformations 1–3, any  $(n, n+1)$ -graph can be changed into one of the graphs  $H_i(p, q)$  ( $i = 1, 2, 3$ ) in Fig. 5, and the Hosoya index increases by these transformations. In order to find the graph with the largest Hosoya index among all  $(n, n+1)$ -graphs, we only need to consider these graphs  $H_i(p, q)$  in Fig. 5.

### 3. The largest Hosoya index of $(n, n+1)$ -graphs

In this section, we first find the graph with the largest Hosoya index in  $\mathcal{A}(p, q)$ , and then find the graphs with the largest Hosoya indices in  $\mathcal{B}(p, q)$  and in  $\mathcal{C}(p, q, r)$ . Finally, we compare the Hosoya indices of these graphs and characterize the extremal graphs.

**Lemma 3.1.** If  $n \geq 6$ ,  $p+q = n+1$ , then  $z(H_1(p, q)) \leq z(H_1(4, n-3))$  with the equality if and only if  $H_1(p, q) \cong H_1(4, n-3)$ .

**Proof.** Let  $m = p = n - q + 1$ . By Eq. (1), we have

$$\begin{aligned} z(H_1(p, q)) &= z(P_{m-1})z(P_{q-1}) + 2z(P_{m-2})z(P_{q-1}) + 2z(P_{m-1})z(P_{q-2}) \\ &= f(m)f(q) + 2f(m-1)f(q) + 2f(m)f(q-1) \\ &= f(m)f(n-m+1) + 2f(m-1)f(n-m+1) + 2f(m)f(n-m) \\ z(H_1(4, n-3)) &= z(P_{n-4})z(P_3) + 2z(P_{n-5})z(P_3) + 2z(P_{n-4})z(P_2) \\ &= f(n-3)f(4) + 2f(n-4)f(4) + 2f(n-3)f(3) \\ &= f(n) + 4f(n-2) \\ &= f(m)f(n-m+1) + f(m-1)f(n-m) + 4f(n-2) \quad (\text{by Lemma 1.1}) \\ z(H_1(4, n-3)) - z(H_1(p, q)) &= f(m-1)f(n-m) + 4f(n-2) - 2f(m-1)f(n-m+1) - 2f(m)f(n-m) \\ &= 2f(n-2) + 2f(n-2) + f(m-1)f(n-m) \\ &\quad - 2f(m-1)f(n-m+1) - 2f(m)f(n-m) \\ &= 2[f(m-1)f(n-m) + f(m-2)f(n-m-1) + f(m)f(n-m-1) \\ &\quad + f(m-1)f(n-m-2) - f(m-1)f(n-m+1) - f(m)f(n-m)] \\ &\quad + f(m-1)f(n-m) \quad (\text{by Lemma 1.1}) \\ &= 2[-f(m-1)f(n-m-1) + f(m-2)f(n-m-1) - f(m)f(n-m-2) \\ &\quad + f(m-1)f(n-m-2)] + f(m-1)f(n-m) \\ &= 2[-f(m-3)f(n-m-1) - f(m-2)f(n-m-2)] + f(m-1)f(n-m) \\ &= 2[-f(m-3)f(n-m-1) - f(m-2)f(n-m-2)] \\ &\quad + f(m-1)f(n-m-1) + f(m-1)f(n-m-2) \\ &= 2[-f(m-3)f(n-m-1) - f(m-2)f(n-m-2)] + f(m-2)f(n-m-1) \\ &\quad + f(m-3)f(n-m-1) + f(m-2)f(n-m-2) + f(m-3)f(n-m-2) \\ &= f(m-4)f(n-m-1) - f(m-4)f(n-m-2) \\ &= f(m-4)f(n-m-3) \geq 0 \end{aligned}$$

with the equality if and only if  $m = 4$  or  $m = n - 3$ , i.e.,  $H_1(p, q) \cong H_1(4, n-3)$ .  $\square$

From Remark 3 and Lemma 3.1, we have immediately:




Fig. 7. The graphs in Lemma 3.3.




Fig. 8. The graphs in Lemma 3.5.

**Theorem 3.2.** Let  $G \in \mathcal{A}(p, q)$ . If  $n \geq 6$ , then  $z(G) \leq z(H_1(4, n-3))$  for all  $p \geq 3$  and  $q \geq 3$ , with the equality if and only if  $G \cong H_1(4, n-3)$ .

If  $n = 5$ , then  $H_1(3, 3)$  is the only graph in  $\mathcal{A}(p, q)$ . So, the graph  $H_1(3, 3)$  or  $H_1(4, n-3)$  is the unique graph with the largest Hosoya index in  $\mathcal{A}(p, q)$  for all  $p \geq 3$  and  $q \geq 3$ .

Next, we show that the Hosoya indices of graphs in  $\mathcal{B}(p, q)$  are less than the largest Hosoya index of graphs in  $\mathcal{A}(p, q)$ .

**Lemma 3.3.** Let  $G$  and  $H$  be the  $(n, n+1)$ -graphs depicted in Fig. 7. Then  $z(G) < z(H)$ .

**Proof.** Let  $M(G)$  and  $M(H)$  be the sets of matchings of  $G$  and  $H$ , respectively. We construct a mapping  $\zeta : M(G) \rightarrow M(H)$ ,  $\forall B \in M(G)$

$$\zeta(B) = \begin{cases} (B - B \cap C_q) \cup \xi_{C_q}(B \cap C_q), & \text{if } vy \in B; \\ B, & \text{otherwise} \end{cases}$$

where  $\xi_{C_q}$  is defined in Lemma 2.3 and  $v$  is taken as  $v_1$ .

Note that  $vz, vx \notin B$  if  $vy \in B$ . The mapping  $\zeta$  is injective. However, there is no  $B \in M(G_1)$  with  $\zeta(B) = \{uy, vx\}$ . So,  $z(G) < z(H)$ .  $\square$

From Remark 3 and Lemmas 3.1 and 3.3, we have:

**Theorem 3.4.** If  $G \in \mathcal{B}(p, q)$ , then  $z(G) < z(H_1(4, n-3))$  for all  $p \geq 3$  and  $q \geq 3$ .

In the following, we give the graph with the largest Hosoya index in  $\mathcal{C}(p, q)$ .

**Lemma 3.5.** Let  $\theta(r, s, t)$  and  $\theta(0, 2, n-4)$  be the  $(n, n+1)$ -graphs depicted in Fig. 8, where  $r + s + t = n-2$ . Then  $z(\theta(r, s, t)) \leq z(\theta(0, 2, n-4))$ , with the equality if and only if  $\theta(r, s, t) \cong \theta(0, 2, n-4)$  or  $\theta(r, s, t) \cong \theta(1, 1, 1)$ .

**Proof.** Let  $G = \theta(r, s, t)$  and  $H = \theta(0, 2, n-4)$ . By Eq. (2), we have

$$\begin{aligned} z(G) &= z(G - \{ux_1\}) + z(G - \{u, x_1\}) \\ &= z(G - \{ux_1, vx_r\}) + z(G - \{ux_1, v, x_r\}) + z(G - \{u, x_1, vx_r\}) + z(G - \{u, x_1, v, x_r\}) \\ &= z(C_{n-r})z(P_r) + 2z(P_{r-1})z(P_{n-r-1}) + z(P_{r-2})z(P_s)z(P_t) \\ &= f(r+1)f(n-r-1) + f(r+1)f(n-r+1) + 2f(r)f(n-r) + f(r-1)f(s+1)f(t+1) \\ z(H) &= z(H - \{uv\}) + z(H - \{u, v\}) \\ &= z(C_n) + z(P_2)z(P_{n-4}) \\ &= f(n-1) + f(n+1) + 2f(n-3). \end{aligned}$$

By Lemma 1.1,

$$\begin{aligned} f(n-r+1) &= f(s+t+3) = f(s+2)f(t+2) + f(s+1)f(t+1) \\ &> 2f(s+1)f(t+1)f(s+1)f(t+1) < \frac{1}{2}f(n-r+1). \end{aligned}$$

If  $r \geq 3$ , then

$$\begin{aligned}
 z(H) - z(G) &= f(n+1) + f(n-1) + 2f(n-3) - f(r+1)f(n-r-1) - f(r+1)f(n-r+1) \\
 &\quad - 2f(r)f(n-r) - f(r-1)f(s+1)f(t+1) \\
 &= f(r+1)f(n-r+1) + f(r)f(n-r) + f(r)f(n-r) + f(r-1)f(n-r-1) \\
 &\quad + 2f(r)f(n-r-2) + 2f(r-1)f(n-r-3) - f(r+1)f(n-r+1) \\
 &\quad - f(r+1)f(n-r-1) - 2f(r)f(n-r) - f(r-1)f(s+1)f(t+1) \quad (\text{by Lemma 1.1}) \\
 &= -f(r-2)f(n-r-1) + 2f(r)f(n-r-2) + 2f(r-1)f(n-r-3) - f(r-1)f(s+1)f(t+1) \\
 &= f(r-2)f(n-r-2) + f(r-2)f(n-r-3) + 2f(r-1)f(n-r-2) \\
 &\quad + 2f(r-3)f(n-r-3) - f(r-1)f(s+1)f(t+1) \\
 &> \frac{1}{2}[2f(r-2)f(n-r-1) + 4f(r-1)f(n-r-2) + 4f(r-3)f(n-r-3) - f(r-1)f(n-r+1)] \\
 &= \frac{1}{2}[2f(r-2)f(n-r-2) + f(r-1)f(n-r-2) + 2f(r-3)f(n-r-3)] \\
 &> 0.
 \end{aligned}$$

If  $r = 1$ , then

$$\begin{aligned}
 z(H) - z(G) &= f(n+1) + f(n-1) + 2f(n-3) - f(n-2) - f(n) - 2f(n-1) \\
 &= f(n-5) \geq 0
 \end{aligned}$$

with the equality if and only if  $n = 5$ .

If  $r = 2$ , then  $s + t = n - 4$  and

$$\begin{aligned}
 z(H) - z(G) &= f(n+1) + f(n-1) + 2f(n-3) - [2f(n-3) + 2f(n-1) + 2f(n-2) + f(s+t)f(t+1)] \\
 &= f(n-3) - f(s+1)f(t+1) \\
 &= f(n-3) - f(s+1)f(n-s-3) \\
 &= f(s)f(n-s-2) + f(s-1)f(n-s-3) - f(s+1)f(n-s-3) \quad (\text{by Lemma 1.1}) \\
 &= f(s)f(n-s-4) \geq 0
 \end{aligned}$$

with the equality if and only if  $s = 0$  or  $t = n - s - 4 = 0$ .  $\square$

From Remark 3 and Lemma 3.5, we have:

**Theorem 3.6.** If  $G \in \mathcal{C}(p, q, r)$ , then  $z(G) \leq z(\theta(0, 2, n-4))$  or  $z(\theta(1, 1, 1))$  for all  $p \geq 3$  and  $q \geq 3$ , with the equality if and only if  $G \cong \theta(0, 2, n-4)$  or  $\theta(1, 1, 1)$ .

Now, in order to find the graph with the largest Hosoya index among all  $(n, n+1)$ -graphs, we only need to compare the Hosoya indices of  $H_1(3, 3)$ ,  $H_1(4, n-3)$ ,  $\theta(0, 2, n-4)$  and  $\theta(1, 1, 1)$ .

$$z(H_1(3, 3)) = 12, \quad z(\theta(1, 1, 1)) = z(\theta(0, 2, 1)) = 13.$$

From the proofs of Lemmas 3.1 and 3.5, we know that

$$z(H_1(4, n-3)) = f(n) + 4f(n-2), \quad z(\theta(0, 2, n-4)) = f(n+1) + f(n-1) + 2f(n-3).$$

Thus  $z(\theta(0, 2, n-4)) - z(H_1(4, n-3)) = 2f(n-5)$ . So, we have:

**Main Theorem.** If  $n \geq 6$ , then  $\theta(0, 2, n-4)$  is the unique graph with the largest Hosoya index among all  $(n, n+1)$ -graphs. If  $n = 5$ , then  $\theta(0, 2, 1)$  and  $\theta(1, 1, 1)$  are the graphs with the largest Hosoya index of all  $(5,6)$ -graphs.

For  $n = 4$ , it is trivial since there is exactly one  $(4,5)$ -graph.

## Acknowledgments

The author would like to thank the referees for their helpful comments and suggestions, which have considerably improved the presentation of the paper.

## References

- [1] H. Hosoya, Topological index, a newly proposed quantity characterizing the topological nature of structural isomers of saturated hydrocarbons, Bull. Chem. Soc. Jpn. 44 (1971) 2332–2339.
- [2] H. Hosoya, Topological index as a common tool for quantum chemistry, statistical mechanics, and graph theory, in: Mathematical and Computational Concepts in Chemistry (Dubrovnik, 1985), in: Ellis Horwood Ser. Math. Appl., Horwood, Chichester, 1986, pp. 110–123.
- [3] H. Deng, S. Chen, The extremal unicyclic graphs with respect to Hosoya index and Merrifield–Simmons index, MATCH Commun. Math. Comput. Chem. 59 (2008) 171–190.

- [4] M. Fischermann, I. Gutman, A. Hoffmann, D. Rautenbach, D. Vidovic, L. Volkmann, Extremal chemical trees, *Z. Naturforsch.* 57a (2002) 49–52.
- [5] M. Fischermann, L. Volkmann, D. Rautenbach, A note on the number of matchings and independent sets in trees, *Discrete Appl. Math.* 145 (3) (2005) 483–489.
- [6] I. Gutman, Extremal hexagonal chains, *J. Math. Chem.* 12 (1993) 197–210.
- [7] I. Gutman, F. Zhang, On the ordering of graphs with respect to their matching numbers, *Discrete Appl. Math.* 15 (1986) 25–33.
- [8] Y. Hou, On acyclic systems with minimal Hosoya index, *Discrete Appl. Math.* 119 (3) (2002) 251–257.
- [9] S. Wagner, Extremal trees with respect to Hosoya index and Merrifield–Simmons index, *MATCH Commun. Math. Comput. Chem.* 57 (2007) 221–233.
- [10] A. Yu, F. Tian, A kind of graphs with minimal Hosoya indices and maximal Merrifield–Simmons indices, *MATCH Commun. Math. Comput. Chem.* 55 (1) (2006) 103–118.
- [11] L.Z. Zhang, The proof of Gutman's conjectures concerning extremal hexagonal chains, *J. Systems Sci. Math. Sci.* 18 (4) (1998) 460–465.
- [12] L. Zhang, F. Tian, Extremal catacondensed benzenoids, *J. Math. Chem.* 34 (2003) 111–122.